

Instructions: Show all work. Some problems will instruct you to complete operations by hand, some can be done in the calculator. To show work on calculator problems, show the commands you used, and the resulting matrices. Give exact answers (yes, that means fractions, square roots and exponentials, and not decimals) unless specifically directed to give a decimal answer. This will require some operations to be done by hand even if not specifically directed to. Be sure to complete all parts of each question.

1. For the vectors 
$$\vec{u} = \begin{bmatrix} 1 \\ 8 \\ -4 \end{bmatrix}$$
,  $\vec{v} = \begin{bmatrix} -1 \\ 7 \\ -2 \end{bmatrix}$ , find the following:

b. A unit vector in the direction of  $\vec{u}$ .

c. 
$$\vec{u} \cdot \vec{v}$$

$$-1+56+8=63$$

Find a vector orthogonal to  $\vec{v}$ .

$$\vec{V} \cdot \vec{N} = -0.176 - 2c = 0$$

$$\dot{y} = 0 \quad \text{then} \\
-0.176 = 0 \\
0.00 = 76 \quad \dot{y} \quad b = 1 \text{ then } n = 7$$

$$a = 76$$

answers may vary.

2. Determine if the polynomials p(t) = t + 6,  $q(t) = t^2 - 3t$  are orthogonal under the inner product  $\langle f|g\rangle = \int_{-1}^1 f(t)g(t)dt$ .

$$\int_{1}^{1} (t+6)(t^{2}-3t)dt =$$

$$\int_{1}^{1} t^{3} - 18t dt = 0 + \int_{1}^{1} 3t^{2} dt$$

= 
$$2\int_{0}^{1} 3t^{2} dt = 2t^{3}\Big|_{0}^{1} = 2$$